




Once you and your doctor have decided that XARELTO® is right for you, XARELTO withMe provides resources and support personalized for you, including cost support options for XARELTO®.

Sign up at XARELTOwithMe.com or call 888-XARELTO (888-927-3586) | Monday–Friday, 8:00 AM–8:00 PM ET

START

XARELTO withMe Trial Offer Free 30-day trial supply for eligible patients



This XARELTO withMe Trial Offer is open to patients who have commercial insurance, government coverage, or no insurance coverage; however, there is no guarantee of continuous accessibility after the program ends.

With the XARELTO withMe Trial Offer, you are able to try XARELTO® at no cost to see if it's right for you. At the conclusion of the program, you and your healthcare provider decide whether to continue treatment.

To be eligible for the XARELTO withMe Trial Offer, you must have been prescribed XARELTO®, except if you are taking XARELTO® 10-mg tablet or 1 mg/mL oral suspension. One (1) use is allowed per lifetime. Terms expire at the end of each calendar year and may change.

To get a XARELTO withMe Trial Offer ID number, talk to your doctor or scan this QR code to receive your Trial Offer information that you can present to your pharmacist with your written prescription for XARELTO® to receive your free trial.



Data rates may apply.

STAY ON XARELTO®

XARELTO withMe Savings Card Eligible patients pay as little as \$10 per fill



There is a limit to savings per fill. Savings may apply to co-pay, co-insurance, or deductible. Your final out-of-pocket cost after the Savings Card is applied will depend on your insurance plan and pharmacy. Not valid for patients using Medicare, Medicaid, or other government-funded programs to pay for their medications. Patients prescribed XARELTO® 10 mg because of a recent non-surgical hospital discharge or because they have recently undergone hip or knee replacement surgery are not eligible. Enrolled patients will receive personalized refill notifications for their XARELTO® prescriptions based on information collected about them and can opt out of refill notifications at any time by calling

888-XARELTO (888-927-3586). Terms expire at the end of each calendar year and may change. Participate without sharing your income information. See Savings Card Program Requirements at XARELTOwithMe.com/SavingsCard.

Sign up at XARELTOwithMe.com or by calling 888-XARELTO (888-927-3586), Monday-Friday, 8:00 AM to 8:00 PM ET.

XARELTO withMe Coverage Gap Support Get help affording your out-of-pocket costs for XARELTO®

If you have insurance covering a portion of XARELTO®, and you pay more than \$89 monthly, XARELTO withMe Coverage Gap Support may be able to help you lower your out-of-pocket costs during a coverage gap—a temporary period where your drug plan limits how much they pay toward your medication. See Coverage Gap Support Program Requirements at XARELTOwithMe.com/CoverageGap.

Sign up at XARELTOwithMe.com or by calling 888-XARELTO (888-927-3586), Monday-Friday, 8:00 AM to 8:00 PM ET.

WHAT IS XARELTO® (rivaroxaban)?

XARELTO® is a prescription medicine used to:

- reduce the risk of stroke and blood clots in adults who have a medical condition called atrial fibrillation that is not caused by a heart valve problem. With atrial fibrillation, part of the heart does not beat the way it should. This can lead to the formation of blood clots, which can travel to the brain, causing a stroke, or to other parts of the body
- treat blood clots in the veins of your legs (deep vein thrombosis or DVT) or lungs (pulmonary embolism or PE)
- reduce the risk of blood clots from happening again in adults who continue to be at risk for DVT or PE after receiving treatment for blood clots for at least 6 months
- help prevent a blood clot in the legs and lungs of adults who have just had hip or knee replacement surgery *continued on page 3*




Once you and your doctor have decided that XARELTO® is right for you, XARELTO withMe provides resources and support personalized for you, including cost support options for XARELTO®.

Sign up at XARELTOwithMe.com or call 888-XARELTO (888-927-3586) | Monday–Friday, 8:00 AM–8:00 PM ET

START

XARELTO withMe Trial Offer Free 30-day trial supply for eligible patients



This XARELTO withMe Trial Offer is open to patients who have commercial insurance, government coverage, or no insurance coverage; however, there is no guarantee of continuous accessibility after the program ends.

With the XARELTO withMe Trial Offer, you are able to try XARELTO® at no cost to see if it's right for you. At the conclusion of the program, you and your healthcare provider decide whether to continue treatment.

To be eligible for the XARELTO withMe Trial Offer, you must have been prescribed XARELTO®, except if you are taking XARELTO® 10-mg tablet or 1 mg/mL oral suspension. One (1) use is allowed per lifetime.

Terms expire at the end of each calendar year and may change.

To get a XARELTO withMe Trial Offer ID number, talk to your doctor or scan this QR code to receive your Trial Offer information that you can present to your pharmacist with your written prescription for XARELTO® to receive your free trial.



Data rates may apply.

STAY ON XARELTO®

XARELTO withMe Coverage Gap Support Get help affording your out-of-pocket costs for XARELTO®

If you have insurance covering a portion of XARELTO®, and you pay more than \$89 monthly, XARELTO withMe Coverage Gap Support may be able to help you lower your out-of-pocket costs during a coverage gap—a temporary period where your drug plan limits how much they pay toward your medication. See Coverage Gap Support Program Requirements at XARELTOwithMe.com/CoverageGap.

Sign up at XARELTOwithMe.com or by calling 888-XARELTO (888-927-3586), Monday-Friday, 8:00 AM to 8:00 PM ET.

XARELTO withMe provides information on other affordability resources that may be available, for example:

- State Health Insurance Programs (SHIPs)
- Medicare Part D Extra Help—Low-Income Subsidy

For more information, call 888-XARELTO (888-927-3586), Monday-Friday, 8:00 AM to 8:00 PM ET or visit JanssenCarePath.com/Xarelto.

Insured patients may be eligible for additional support from Janssen

Patient assistance is available if you have commercial, employer-sponsored, or government coverage that does not fully meet your needs. You may be eligible to receive your Janssen medication free of charge for up to one year. You must meet the eligibility and income requirements for the Janssen Patient Assistance Program. See terms and conditions at PatientAssistanceInfo.com.

XARELTO withMe is limited to education about XARELTO®, its administration, and/or the condition it treats. It is not intended to provide medical advice, replace a treatment plan you receive from your doctor or nurse, or serve as a reason for you to start or stay on treatment.

WHAT IS XARELTO® (rivaroxaban)? (continued)

- help prevent blood clots in certain adults hospitalized for an acute illness and after discharge, who are at risk of getting blood clots because of the loss of or decreased ability to move around (mobility) and other risks for getting blood clots, and who do not have a high risk of bleeding

XARELTO® is used with low dose aspirin to:

- reduce the risk of serious heart problems, heart attack and stroke in adults with coronary artery disease (a condition where the blood supply to the heart is reduced or blocked)
- reduce the risk of a sudden decrease in blood flow to the legs, major amputation, serious heart problems or stroke in adults with peripheral artery disease (a condition where the blood flow to the legs is reduced) and includes adults who have recently had a procedure to improve blood flow to the legs

IMPORTANT SAFETY INFORMATION

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT XARELTO®?

XARELTO® may cause serious side effects, including:

- **Increased risk of blood clots if you stop taking XARELTO®.** People with atrial fibrillation (an irregular heart beat) that is not caused by a heart valve problem (nonvalvular) are at an increased risk of forming a blood clot in the heart, which can travel to the brain, causing a stroke, or to other parts of the body. XARELTO® lowers your chance of having a stroke by helping to prevent clots from forming. If you stop taking XARELTO®, you may have increased risk of forming a clot in your blood.

Do not stop taking XARELTO® without talking to the doctor who prescribes it for you.

Stopping XARELTO® increases your risk of having a stroke. If you have to stop taking XARELTO®, your doctor may prescribe another blood thinner medicine to prevent a blood clot from forming.

- **Increased risk of bleeding.** XARELTO® can cause bleeding which can be serious and may lead to death. This is because XARELTO® is a blood thinner medicine (anticoagulant) that lowers blood clotting. During treatment with XARELTO® you are likely to bruise more easily, and it may take longer for bleeding to stop. You may be at higher risk of bleeding if you take XARELTO® and have certain other medical problems.

You may have a higher risk of bleeding if you take XARELTO® and take other medicines that increase your risk of bleeding, including:

- Aspirin or aspirin-containing products
- Long-term (chronic) use of non-steroidal anti-inflammatory drugs (NSAIDs)
- Warfarin sodium (Coumadin®, Jantoven®)
- Any medicine that contains heparin
- Clopidogrel (Plavix®)
- Selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs)
- Other medicines to prevent or treat blood clots

Please see Important Safety Information continued on pages 4 and 5, and read the full [Prescribing Information](#), including [Boxed Warnings](#), and [Medication Guide](#) for XARELTO®, and discuss any questions you have with your doctor.

XARELTO® is used in children to:

- treat blood clots or reduce the risk of blood clots from happening again in children from birth to less than 18 years, after receiving at least 5 days of treatment with injectable or intravenous medicines used to treat blood clots
- help prevent blood clots in children 2 years and older with congenital heart disease after the Fontan procedure

XARELTO® was not studied and is not recommended in children less than 6 months of age who:

- were less than 37 weeks of growth (gestation) at birth
- had less than 10 days of oral feeding, **or**
- had a body weight of less than 5.7 pounds (2.6 kg)

Tell your doctor if you take any of these medicines. Ask your doctor or pharmacist if you are not sure if your medicine is one listed above.

Call your doctor or get medical help right away if you or your child develop any of these signs or symptoms of bleeding:

- Unexpected bleeding or bleeding that lasts a long time, such as:
 - Nosebleeds that happen often
 - Unusual bleeding from gums
 - Menstrual bleeding that is heavier than normal, or vaginal bleeding
- Bleeding that is severe or you cannot control
- Red, pink, or brown urine
- Bright red or black stools (looks like tar)
- Cough up blood or blood clots
- Vomit blood or your vomit looks like “coffee grounds”
- Headaches, feeling dizzy or weak
- Pain, swelling, or new drainage at wound sites
- **Spinal or epidural blood clots (hematoma).** People who take a blood thinner medicine (anticoagulant) like XARELTO®, and have medicine injected into their spinal and epidural area, or have a spinal puncture, have a risk of forming a blood clot that can cause long-term or permanent loss of the ability to move (paralysis). Your risk of developing a spinal or epidural blood clot is higher if:
 - A thin tube called an epidural catheter is placed in your back to give you certain medicine
 - You take NSAIDs or a medicine to prevent blood from clotting
 - You have a history of difficult or repeated epidural or spinal punctures
 - You have a history of problems with your spine or have had surgery on your spine

If you take XARELTO® and receive spinal anesthesia or have a spinal puncture, your doctor should watch you closely for symptoms of spinal or epidural blood clots.

Tell your doctor right away if you have:

- back pain

continued next page

IMPORTANT SAFETY INFORMATION (continued)

- tingling
- numbness
- muscle weakness (especially in your legs and feet)
- or loss of control of the bowels or bladder (incontinence)

XARELTO® is not for use in people with artificial heart valves.

XARELTO® is not for use in people with antiphospholipid syndrome (APS), especially with positive triple antibody testing.

Do not take XARELTO® if you or your child:

- Currently have certain types of abnormal bleeding. Talk to your doctor before taking XARELTO® if you currently have unusual bleeding.
- Are allergic to rivaroxaban or any of the ingredients of XARELTO®.

Before taking XARELTO®, tell your doctor about all your medical conditions, including if you or your child:

- Have ever had bleeding problems
- Have liver or kidney problems
- Have antiphospholipid syndrome (APS)
- Are pregnant or plan to become pregnant. It is not known if XARELTO® will harm your unborn baby.
 - **Tell your doctor** right away if you become pregnant during treatment with XARELTO®. Taking XARELTO® while you are pregnant may increase the risk of bleeding in you or in your unborn baby.
 - Females who are able to become pregnant: Talk with your doctor about pregnancy planning during treatment with XARELTO®. Talk with your doctor about your risk for severe uterine bleeding if you are treated with blood thinner medicines, including XARELTO®.
 - If you take XARELTO® during pregnancy, **tell your doctor** right away if you have any signs or symptoms of bleeding or blood loss. **See “What is the most important information I should know about XARELTO®?” for signs and symptoms of bleeding.**
- Are breastfeeding or plan to breastfeed. XARELTO® may pass into your breast milk. Talk to your doctor about the best way to feed your baby during treatment with XARELTO®.

Tell all of your doctors and dentists that you or your child are taking XARELTO®. They should talk to the doctor who prescribed XARELTO® for you before you have any surgery, medical or dental procedure.

Tell your doctor about all the medicines you or your child take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some of your other medicines may affect the way XARELTO® works, causing side effects. Certain medicines may increase your risk of bleeding. **See “What is the most important information I should know about XARELTO®?”**

HOW SHOULD I TAKE XARELTO®?

- Take XARELTO® exactly as prescribed by your doctor.
- **Do not change your dose or stop taking XARELTO® unless your doctor tells you to.** Your doctor may change your dose if needed.
- Your doctor will decide how long you should take XARELTO®.
- XARELTO® may need to be stopped for one or more days before any surgery or medical or dental procedure. Your doctor will tell you when to stop taking XARELTO® and when to start taking XARELTO® again after your surgery or procedure.
- If you need to stop taking XARELTO® for any reason, talk to the doctor who prescribed XARELTO® to you to find out when you should stop taking it. Do not stop taking XARELTO® without first talking to the doctor who prescribes it to you.
- If you have difficulty swallowing XARELTO® tablets whole, talk to your doctor about other ways to take XARELTO®.
- Do not run out of XARELTO®. Refill your prescription of XARELTO® before you run out. When leaving the hospital following a hip or knee replacement, be sure that you will have XARELTO® available to avoid missing any doses.
- If you take too much XARELTO®, go to the nearest hospital emergency room or call your doctor right away.

If you take XARELTO® for:

• Atrial Fibrillation that is not caused by a heart valve problem:

- Take XARELTO® **1 time a day with your evening meal.**
- If you miss a dose of XARELTO®, take it as soon as you remember on the same day. Take your next dose at your regularly scheduled time.

• Blood clots in the veins of your legs or lungs:

- Take XARELTO® **1 or 2 times a day** as prescribed by your doctor.
- For the **10-mg dose**, XARELTO® may be taken with or without food.
- For the **15-mg and 20-mg doses**, take XARELTO® **with food at the same time each day.**
- If you miss a dose:
 - **If you take the 15-mg dose of XARELTO® 2 times a day (a total of 30 mg of XARELTO® in 1 day):** Take XARELTO® as soon as you remember on the same day. You may take 2 doses at the same time to make up for the missed dose. Take your next dose at your regularly scheduled time.
 - **If you take XARELTO® 1 time a day:** Take XARELTO® as soon as you remember on the same day. Take your next dose at your regularly scheduled time.

• Hip or knee replacement surgery:

- Take XARELTO® 1 time a day with or without food.

continued next page

Please see Important Safety Information continued on page 5, and read the full [Prescribing Information](#), including [Boxed Warnings](#), and [Medication Guide for XARELTO®](#), and discuss any questions you have with your doctor.

IMPORTANT SAFETY INFORMATION (continued)

- If you miss a dose of XARELTO®, take it as soon as you remember on the same day. Take your next dose at your regularly scheduled time.
- **Blood clots in people hospitalized for an acute illness:**
 - Take XARELTO® 1 time a day, with or without food, while you are in the hospital and after you are discharged as prescribed by your doctor.
 - If you miss a dose of XARELTO®, take it as soon as you remember on the same day. Take your next dose at your regularly scheduled time.
- **Reducing the risk of serious heart problems, heart attack and stroke in coronary artery disease:**
 - Take XARELTO® 2.5 mg 2 times a day with or without food.
 - If you miss a dose of XARELTO®, take your next dose at your regularly scheduled time.
 - Take aspirin 75 to 100 mg once daily as instructed by your doctor.
- **Reducing the risk of a sudden decrease in blood flow to the legs, major amputation, serious heart problems or stroke in people with peripheral artery disease, including those who have recently had a procedure to improve blood flow to the legs:**
 - Take XARELTO® 2.5 mg 2 times a day with or without food.
 - If you miss a dose of XARELTO®, take your next dose at your regularly scheduled time.
 - Take aspirin 75 to 100 mg once daily as instructed by your doctor.

For children who take XARELTO®:

- The dose of XARELTO® depends on your child’s body weight and will be calculated by your child’s doctor. Your child’s doctor will tell you if XARELTO® can be given to your child with or without food.
- The adult caregiver should give the dose.
- If your child is taking the tablet, the tablet should be taken whole and should not be split in an attempt to provide a lower dose of XARELTO®.
- If your child is taking the oral suspension, use the syringes provided in the original carton. The suspension will be prepared by the pharmacy. See the **Instructions for Use** included in the carton on how to properly give a dose of XARELTO® oral suspension to your child.
- Do not switch between the XARELTO® oral suspension or tablet without first talking to your doctor.
- If your child vomits or spits up:
 - right after or within 30 minutes of taking the oral suspension, give a new full dose.
 - more than 30 minutes after taking the oral suspension, do not give the dose again. Give the next dose at the regularly scheduled time.
 - if vomiting or spitting up persists, contact your child’s doctor right away.

- If your child misses a dose:
 - If your child is taking XARELTO® 1 time a day, give the dose as soon as you remember on the same day. If this is not possible, skip this dose and give the next dose at the regularly scheduled time.
 - If your child is taking XARELTO® 2 times a day, give the missed morning dose as soon as you remember. You may give the missed morning dose together with the evening dose. However, a missed evening dose can only be taken in the same evening.
 - If your child is taking XARELTO® 3 times a day, skip the missed dose and give the next dose at the regularly scheduled time.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF XARELTO®?

XARELTO® may cause serious side effects:

- See **“What is the most important information I should know about XARELTO®?”**

The most common side effect of XARELTO® in adults was bleeding.

The most common side effects of XARELTO® in children include:

- bleeding
- cough
- vomiting
- inflamed stomach and gut

Call your doctor for medical advice about side effects. **You may report side effects to FDA at 1-800-FDA-1088.**

You may also report side effects to Janssen Pharmaceuticals, Inc., at 1-800-JANSSEN (1-800-526-7736).

Please read the full Prescribing Information, including Boxed Warnings, and Medication Guide for XARELTO®, and discuss any questions you have with your doctor

Trademarks are those of their respective owners.

XARELTO® is licensed from Bayer HealthCare AG, 51368 Leverkusen, Germany.